



King's Research Portal

DOI:

[10.1071/MF16373](https://doi.org/10.1071/MF16373)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Baker, K., Chadwick, M. A., McGill, R. A. R., Wahab, R. A., & Kahar, R. (2017). Macroinvertebrate trophic structure on waterfalls in Borneo. *MARINE AND FRESHWATER RESEARCH*, 68(11), 2061-2069. <https://doi.org/10.1071/MF16373>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Title: Macroinvertebrate trophic structure on waterfalls in Borneo

Short title: Macroinvertebrate trophic structure on Waterfalls

Authors: Kate Baker^{1*}, Michael A. Chadwick¹, Rona A.R. McGill², Rodzay A. Wahab³ and Rafhiah Kahar³

¹ Department of Geography, King's College London, UK

² NERC Life Sciences Mass Spectrometry Facility (LSMSF), Scottish Universities Environmental Research Centre, Rankine Avenue, East Kilbride, Glasgow G75 0QF

³ Universiti Brunei Darussalam, Bandar Seri Begawan, Brunei Darussalam

*Corresponding author; e-mail: katedotbaker@gmail.com and Tel: +447748027326.

Abstract

Waterfalls have unique physical characteristics and harbour highly specialized macroinvertebrate communities, but have been the subject of very few ecological studies. There are no previous studies of trophic structure of waterfall assemblages. It was hypothesized that due to the steep gradient of waterfalls and low retention of terrestrial based resources, the abundant basal food resources would be periphyton. In addition, due to the frequent scouring flood events, it was predicted that periphyton would also be a significant source of food for filter feeders. Waterfalls in the Ulu Temburong National Park in Brunei Darussalam (northern Borneo) were used as this case study. Methods included stable isotope analysis (SIA; $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of leaf litter and periphyton) and gut contents analysis (GCA) of the most the abundant macroinvertebrates. $\delta^{15}\text{N}$ values ranging from -1.9‰ to 5.5‰, literature suggests that this indicates there were herbivores (Heptageniidae and Blephariceridae), omnivores (Simuliidae and Hydropsychidae) and predators (Buccinidae) living on the waterfalls. Apart from Buccinidae, all the taxa had $\delta^{13}\text{C}$ signatures ranging from -33‰ to -26‰, with a high dependence on periphyton, which is like other tropical stream biotopes. This study does suggest that despite the scouring velocities, waterfalls support animals with a range of diets, based on grazing/scraping, filter feeding and predation.

Introduction

Waterfalls have received limited attention with very few published papers to date (Rackemann *et al.* 2013; Clayton and Pearson, 2016). Research conducted on waterfalls in the tropics have generally focused on them acting as a longitudinal barrier for fish movement, creating an upstream refuge for taxa including shrimp and tadpoles (Baker *et al.* 2016; Covich *et al.* 2009; Hein and Crowl, 2010; Torrente-Vilara *et al.* 2011; Kano *et al.* 2012), with very little information on the waterfall assemblages themselves (Clayton and Pearson, 2016; Rackemann *et al.* 2013 are exceptions). This lack of research may be explained by several factors including waterfalls being perceived to be biological dead zones, too complex to sample effectively and being inherently dangerous environments due to fast flows and sheer drops (Rackemann *et al.* 2013).

Waterfalls are defined by their bedrock substrate and fast flows (Newson and Newson, 2000) and are unique compared to other biotopes because of their sheer shallow flows and lack of connectivity to the hyporheic zones. In addition, these biotopes are formed and maintained by erosive processes that occur over geologic periods creating stable conditions, which are relatively uniform across a range of discharges (Clayton and Pearson, 2016). In comparison, other biotopes such as pools and riffles change daily, frequently re-shaped by erosion and deposition events, whereas waterfalls are more stable (Clayton and Pearson, 2016).

Although waterfalls have stable substrate, resident taxa have to be specialized to live in an environment with relatively fast velocities and no hyporheic zone (Clayton and Pearson, 2016; Hart and Finelli, 1999). Animals need morphological adaptations,

including hooks, suckers and modified legs, to enable them to move around the rock without being washed away or to fit into cracks and crevices (Clayton and Pearson, 2016). For example, some odonates in a Malaysian stream inhabited trailing plants and accumulated debris irrespective of water velocity, assisted by their body spines, which aided attachment (Furtado 1969). In addition to the high velocity areas of the waterfalls, there are some benign zones including the waterfall spray, these patches are usually colonised by snails.

Taxa adapted to living on waterfalls benefit from the absence of fish and shrimp, which have been found to be important in determining insect assemblage structure (March *et al.* 2002; March and Pringle, 2003; Ramirez and Hernandez-Cruz, 2004). In streams, fish and shrimp typically impact insects via predation and competition for food resources, breaking up and consuming leaf litter and grazing periphyton (Pringle and Hamazaki, 1997; Crowl *et al.* 2001; Flecker and Taylor, 2004; Ramirez and Hernandez-Cruz, 2004; Coughlan *et al.* 2010). In addition, fish and shrimp can occur in high densities, creating competition for space with invertebrates. Waterfalls are therefore unique biotopes for invertebrates as they can reside on the substrate with minimal top-down ecological disturbance. It is therefore predicted that the difference in community structure on waterfalls, will create a distinctive food web compared to other tropical biotopes (e.g. Brito *et al.* 2006; Li and Dudgeon, 2008; Coat *et al.* 2009).

Food web studies using stable isotope analysis in tropical streams have focused on pool and riffle biotopes dominated by shrimp and fish (for example see: Brito *et al.* 2006; Li and Dudgeon, 2008; Coat *et al.* 2009). Some of these studies have found invertebrates in the tropics to depend more on algal based food compared to leaf litter (Brito *et al.* 2006; Lau *et al.* 2009). Even in shaded reaches, some tropical food webs studies (e.g. in the Neotropics) have been found to be mainly algae based (March and

Pringle, 2003; Brito *et al.* 2006). Gaps in the canopy often occur near to waterfall, including upstream, below and along the waterfall, as a result of the high rates of erosion below the waterfalls, which create wide plunge pools (Odland *et al.* 1991). This canopy gap enables more light to reach the waterfall substrate promoting periphyton growth. In contrast, standing stocks of benthic organic matter tend to be low on waterfalls because of fast flows, channel gradient and the lack of retentive structure. Although waterfalls are discrete units they are not separated from the river continuum with resident filter feeders, such as Simuliidae and Hydropsychidae, dependent on seston and organic matter from upstream. However, the strength of upstream-downstream linkages have been debated with Huryn *et al.* (2002) suggesting that local factors are still more of a determinant of trophic resources.

This study examined macroinvertebrate trophic structure on waterfalls in Ulu Temburong Park, which has extensive pristine tropical rainforest with free flowing rivers (Sheldon, 2011). The study combines complementary techniques of gut contents analysis (GCA) and stable isotope analysis (SIA), to enable a more complete characterization of waterfall trophic structure. SIA is useful for tracing energy flows through food webs but gives poor taxonomic resolution (Fry, 2007; Grey *et al.* 2001). In contrast, GCA can provide higher taxonomic resolution but provides only a snapshot of ingestion rather than assimilation (Layer *et al.* 2013). It was hypothesized that due to the steep gradient of waterfalls and low retention of terrestrial based resources, the predominant basal food resources of macroinvertebrates would be periphyton. In addition, due to the frequent scouring flood events, it was predicted that periphyton would also be a significant source of food for filter feeders.

Methods

Study Sites

Research was conducted at the Kuala Belalong Field Studies Centre (KBFS) of UBD (Universiti Brunei Darussalam) in Ulu Temburong National Park in Brunei Darussalam. The landscape of Ulu Temburong National Park (550 km²) changes over a relatively short distance (< 35 km) from steep mountainous reaches (913 m a.s.l. at Bukit Belalong) to placid lowlands (KBFS 30 m a.s.l.) (Dykes, 1994). This creates a geomorphic template for many waterfalls and cascades, therefore a suitable region for this study (Fig. 1).

Brunei is in the tropics and weakly influenced by the South East Asian monsoon; however, weather in Ulu Temburong is highly variable as a result of localised storms (Dykes, 1996). Sampling took place in June and July (2014) when rainfall and stream discharges are generally low making access to the waterfalls safe (Cranbrook and Edwards, 1994). Individual waterfalls on twelve tributaries of Sungai Temburong or Sungai Belalong were investigated (Fig. 1). Waterfalls selected to be sampled were the first upstream after the confluence with the main rivers.

Geomorphic Measurements and Functional Habitats

We measured width and length of the waterfalls, except where waterfalls were very steep, in which case their length was estimated. The presence and distribution of functional habitats (e.g. moss, leaf litter, wood debris) were recorded (for an explanation of functional habitats see Harper *et al.* 1995). Descriptions of substrate heterogeneity and photos of the waterfalls were taken to enable qualitative image analysis of waterfall substrate complexity.

165 *Macroinvertebrate trophic structure*

166 The common and widely distributed taxa on the waterfalls, including Heptageniidae,
167 Blephariceridae, Simuliidae, Hydropsychidae, and Buccinidae, were picked directly off
168 the waterfall substrate, either by hand or with forceps. Sampling was conducted over
169 the length of the waterfall, except on the larger waterfalls where it was unsafe to sample
170 in the middle. Trophic structure was then evaluated using gut contents analysis (GCA)
171 and stable isotope analysis (SIA). Species richness within these families in streams
172 surrounding KBFSC is uncertain. However, to date, three genera of Heptageniidae
173 (*Componeuria*, *Thalerosphyrus* and *Afronurus*), one genus of Blephariceridae
174 (*Blepharicera*), six genera of Hydropsychidae (*Hydromanicus*, *Macrostemum*,
175 *Hydropsyche*, *Cheumatopsyche*, *Potamyia* and *Polymorphanisus*) and one genus of
176 Buccinidae (*Clea*) have been recorded (Baker et al., in press). To date *Clea* is the only
177 freshwater genus from the Buccinidae family. The level of diversity within Simuliidae
178 is unknown in the streams of Ulu Temburong National Park.

179
180 *Gut Contents Analysis:* Gut contents analysis (GCA) has been used frequently to
181 investigate benthic macroinvertebrate diet (Rosi-Marshall and Wallace, 2002; Li and
182 Dudgeon, 2008). Due to the remote location, only three individuals of each of the taxon
183 were collected for GCA. Macroinvertebrates were preserved in 70% ethanol. In the
184 laboratory, foreguts were removed from the individuals under a dissecting microscope
185 and the contents mounted in Euparal on a microscope slide (Layer *et al.* 2012). The
186 contents were identified at 400-1,000x magnification and separated into five categories:
187 coarse particulate organic matter (CPOM) including leaf litter and wood, algae
188 (diatoms, green alga), fungal hyphae and conidia, animal tissue and amorphous detritus

identified by the lack of any defined cellular structure; from Yule *et al.* (2010).

Percentage of gut contents was then estimated by eye.

Stable Isotope Analysis: Stable isotopes have been widely used to evaluate energy flow and trophic structure in a range of food webs (Peterson and Fry, 1987). Stable carbon and nitrogen isotope ratios are a useful tool to calculate what has been assimilated by animals and to determine the relative importance of basal food resources in the food web. Carbon isotope ratios are effective in distinguishing between autochthonous (aquatic) and allochthonous (terrestrial) food sources, with the former having higher $\delta^{13}\text{C}$ values (Fry, 1991; Hershey and Peterson, 1996). Nitrogen isotopes have been used to provide information about trophic levels, with $\delta^{15}\text{N}$ increasing with each trophic level.

Invertebrates were collected for stable isotope analyses using the same methods as gut contents analysis (see above). From 3-50 individuals per taxon were selected to ensure enough sample material for analysis. All taxa were left in stream water for 12 hours to promote gut clearance. Putative food sources (leaf litter and periphyton rock scrubs) were also taken at each waterfall. All samples were dried at 70 degrees Celsius for a 24hour period before being transported back to the UK. Samples were weighed to approximately 0.5mg and loaded into 5x7mm tin capsules. Leaf litter was ground into a fine powder. Initial analysis found that three replicates were sufficient for macroinvertebrates and leaf litter, and due to the increased variability, five replicates were required for the periphyton.

Measurements were carried out at the NERC Life Sciences Mass Spectrometry Facility (LSMSF) in East Kilbride, using an Elementar Pyrocube elemental analyser coupled with a Thermo Fisher Scientific Delta Plus XP mass spectrometer. Laboratory standards Fluka gelatin, Sigma alanine and Sigma glycine (Sigma-Aldrich Company

Ltd, Gillingham, UK) were repeated with every 10 samples and were used to correct for linearity and instrument drift over a 22 hour analytical run. The isotope ratios for the lab standards were determined relative to a range of International standards from IAEA (Vienna, Austria) and USGS (Reston, VA, USA). Isotope ratios are expressed in the δ (delta) notation in parts per million (‰): $\delta X = [(R_{\text{sample}}/R_{\text{standard}}) - 1]$ where $X = ^{15}\text{N}$ or ^{13}C and R = the ratio of $^{15}\text{N}/^{14}\text{N}$ or $^{13}\text{C}/^{12}\text{C}$ isotopes in a given sample relative to AIR for nitrogen and PDB for carbon. The analytical precision for carbon and nitrogen isotope measurements was better than 0.3‰.

Data Analysis: Analysis of variance (ANOVA) was used to assess differences in stable isotope compositions for food sources and consumers. It was also used to assess differences among all the waterfalls. Tukey Post Hoc test were used after the ANOVA to examine specific differences (Thomas *et al.* 2013). The statistical computing environment R (R Core Team 2013) was used to conduct the ANOVAs. The focus of the study was to examine allochthonous (leaf litter) and autochthonous (periphyton) food sources on the waterfalls. As not all food sources were sampled (e.g. moss, seston, fungi), a mixing model was not applied to the data.

Results

Geomorphic Measurements and Functional Habitats

The twelve waterfalls had a range of average slope angles, from 16° (waterfall 11) to 53° (waterfall 1) and lengths of the waterfalls were from 4.5m (waterfall 11) to 20.4m (waterfall 8-Table 1). All submerged rock surfaces were covered in periphyton with at least a few leaves present. Half of the waterfalls had wood debris, which varied in size from twigs to large tree trunks. At waterfalls 1, 4 and 7, large tree trunks became

Commented [BK1]: that is not parts per million, it is parts per thousand (sometimes expressed as parts per mil, mil being short for the Latin word for thousand)

lodged in between the bedrock and the narrow bankful widths. Our observations in the field and subsequent qualitative image analysis indicated that waterfalls varied in substrate complexity. Some of the waterfalls consisted of smooth rock while others were more complex with a mix of ridges (millimetres to centimetres), ledges and undercutting.

Macroinvertebrate trophic structure

The sampled macroinvertebrates included Simuliidae, Hydropsychidae, Blephariceridae, Heptageniidae and Buccinidae. At waterfalls 8 and 11 there were no Blephariceridae and no Buccinidae at waterfall 8.

Gut Contents Analysis; Amorphous detritus was most abundant material in macroinvertebrate guts, ranging from ~45% to ~100% (Fig. 2). CPOM was the next most abundant, ranging from 0 to 30% while algae varied from 0 to ~20% (Fig. 2). Animal tissue varied from 0 to ~10% and the least abundant matter in the macroinvertebrate guts was fungal hyphae and conidia varying from 0 to 5% (Fig. 2). Blephariceridae contained 100% amorphous detritus. Simuliidae contained a high percentage of amorphous detritus (~95%), but in addition included CPOM, algae, and in one individual an insect head (Waterfall 9; Fig. 2). Heptageniidae contained a high percentage of amorphous detritus, but also contained algae and fungal hyphae and conidia (Fig. 2). Hydropsychidae had a more even mix of all observed food types (Fig. 2).

Stable isotopic composition of food sources: Periphyton had less depleted ^{13}C values ($\delta^{13}\text{C}$ periphyton = -30.06 ± 2.54 ‰) compared to leaf litter ($\delta^{13}\text{C}$ leaf litter = $-32.97 \pm$

1.58 ‰), making it possible to distinguish between these basal food resources (one-way ANOVA, $F_{1, 83}=37.7$, $p<0.01$; Table 2; Appendix 1). The $\delta^{15}\text{N}$ (‰) values for periphyton (1.29 ± 0.78 ‰) and leaf litter (-0.91 ± 0.99 ‰) were also distinct (one-way ANOVA, $F_{1, 83}=142.6$, $p<0.01$), with periphyton having significantly higher $\delta^{15}\text{N}$ values than leaf litter. The C: N values were higher for the leaf litter (49.27) than for periphyton (5.37). The lower ratio indicates a higher proportion of nitrogen in the sample suggesting higher protein digestibility and therefore higher nutritional value to the consumers (Giller and Malmqvist, 1998; Table 2).

Stable isotopic compositions of consumers: The $\delta^{13}\text{C}$ values for Heptageniidae at the 12 waterfalls varied from -35.51 ‰ to -26.83 ‰, while Blephariceridae ranged from -35.68 ‰ to -27.41 ‰ (Table 2; Fig. 3; Appendix 1). Simuliidae varied from -29.29 ‰ to -27.33 ‰ and Hydropsychidae $\delta^{13}\text{C}$ values ranged from -31.12 ‰ to -27.2‰. Buccinidae ranged from -28.03 ‰ to -24.04 ‰ (Table 2; Fig. 3). There was a statistical difference between the $\delta^{13}\text{C}$ values of all consumers (one-way ANOVA, $F_{4, 151}=55.5$, $p<0.01$), but the Tukey post hoc test showed no difference among Heptageniidae and Blephariceridae. Mean $\delta^{15}\text{N}$ values of Heptageniidae ranged from a -0.56 ‰ to 2.51 ‰, Blephariceridae varied from 0.51 ‰ to 2.64 ‰ and Simuliidae varied from 1.65 ‰ to 3.04 ‰ (Table 2). Hydropsychidae ranged from 2.26 ‰ to 4.50 ‰ and Buccinidae ranged from 4.12 ‰ to 6.69 ‰ (Table 2). There was a statistical difference between the consumers in $\delta^{15}\text{N}$ values (one-way ANOVA, $F_{4, 151}=275.7$, $p<0.01$). However, the Tukey post hoc test showed no difference in $\delta^{15}\text{N}$ values between Heptageniidae and Blephariceridae ($p=0.34$), and between Simuliidae and Hydropsychidae ($p=0.8$).

Potential food sources of consumers using GCA and SIA: Most Heptageniidae had $\delta^{13}\text{C}$ values that corresponded to periphyton (Table 2; Fig. 3). However, at waterfall 7,

289 Heptageniidae had a $\delta^{13}\text{C}$ value of approximately -27 ‰ and at waterfall 10 the $\delta^{13}\text{C}$
 290 values were very depleted at approximately -35 ‰ (Table 2; Fig. 3). GCA found
 291 Heptageniidae consumed CPOM, algae and, fungal hyphae and conidia, the latter food
 292 source may explain the 'unknown' $\delta^{13}\text{C}$ values. At most waterfalls Blephariceridae
 293 matched periphyton and leaf litter; however, at waterfall 10 $\delta^{13}\text{C}$ values were depleted
 294 with $\delta^{13}\text{C}$ values less than -35 ‰. Due to the small size of the Blephariceridae, the GCA
 295 could not be used to show any further resolution, with gut material classed as
 296 amorphous detritus. Generally Simuliidae $\delta^{13}\text{C}$ values matched periphyton, although,
 297 there were some Simuliidae at waterfall 11 that had $\delta^{13}\text{C}$ value of approximately -27
 298 ‰, values that are on the edge of the periphyton $\delta^{13}\text{C}$ (Fig. 3). The GCA did not provide
 299 any further resolution (Fig. 2). Hydropsychidae $\delta^{13}\text{C}$ values were similar to periphyton,
 300 along with another food source with $\delta^{13}\text{C}$ of approximately -27 ‰ at waterfall 8 and
 301 11. GCA of Hydropsychidae contained animal tissue, fungal hyphae and conidia,
 302 CPOM and algae. Buccinidae $\delta^{13}\text{C}$ values did not match periphyton or leaf litter apart
 303 from at waterfall 9 where $\delta^{13}\text{C}$ values corresponded to periphyton. Buccinidae gut
 304 contents was amorphous therefore not possible to differentiate food types.
 305 Range of $\delta^{15}\text{N}$ values indicate that macroinvertebrates on the 12 waterfalls fed at
 306 different trophic levels (Fig. 4). Heptageniidae had the most depleted ^{15}N with a value
 307 of 1.06 ± 0.77 ‰ and Blephariceridae had a $\delta^{15}\text{N}$ value of 1.50 ± 0.68 ‰ (Table 2; Fig.
 308 4). Simuliidae and Hydropsychidae $\delta^{15}\text{N}$ values were higher, Simuliidae had average
 309 $\delta^{15}\text{N}$ value of 2.16 ± 0.36 ‰ and Hydropsychidae $\delta^{15}\text{N}$ value was 3.07 ± 0.46 ‰ (Table
 310 2; Fig. 3). This indicates that some of their diet was composed of animal tissue. GCA
 311 showed that animal tissue was present in the stomach contents of Simuliidae, although
 312 only at waterfall 9. Hydropsychidae had animal tissue in gut contents for half of the
 313 waterfalls including 2, 3, 7, 8, 9 and 12. Whilst Buccinidae was distinctly predacious,

314 having the most enriched ^{15}N with values of 5.48 ± 0.70 ‰ (Table 2; Fig. 3). These
315 results suggest that there were at least two trophic levels, with Heptageniidae and
316 Blephariceridae as primary consumers and Buccinidae as a secondary consumer.
317 Simuliidae and Hydropsychidae were in the middle, with some taxa primary consumers
318 and others secondary (Fig. 4).

319 Discussion

320 **There are few ecological studies of waterfalls** (Rackemann *et al.* 2013; Clayton
321 and Pearson, 2016) and we are unaware of any published papers on waterfall food webs.
322 Waterfalls are unique biotopes, lacking a hyporheic zone and a water column of
323 sufficient depth for most fish. Therefore the normally abundant herbivorous fish and
324 shrimp are absent, leaving highly specialized macroinvertebrates to dominate the
325 waterfalls (Clayton and Pearson, 2016; Hart and Finelli, 1999). This study, using SIA
326 and GCA, has established a trophic structure for waterfalls in pristine catchments in
327 northern Borneo showing there are at least two, and potentially three, trophic levels –
328 detritivore/herbivore (Heptageniidae and Blephariceridae), omnivore (Simuliidae and
329 Hydropsychidae) and predator (Buccinidae). There was some variation in carbon and
330 nitrogen isotopic values of macroinvertebrates among the waterfalls, which is likely
331 linked to variations in functional habitats including both quantity and quality. Further
332 research relating functional habitats and waterfall geomorphology to the trophic
333 ecology of waterfall resident taxa would be required to elucidate these patterns.

334 The isotopic values ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) of basal food sources were distinct for
335 periphyton and leaf litter (Table 2), making it possible to use them as indicators to
336 distinguish between autochthonous and allochthonous basal food resources. The $\delta^{13}\text{C}$
337 values for leaf litter averaged -32.97 ‰, lower compared to other values reported from
338

the tropics. For example, Huang *et al* (2007) recorded CPOM from streams in Taiwan with an average of -30.3 ‰ and values from Hong Kong averaged -31.4 ‰ (Mantel *et al.* 2004). This range in $\delta^{13}\text{C}$ values can be likely attributed to the high tree diversity in tropical rainforests – for example in Ulu Temburong national park there are approximately 250 species of trees (Cranbrook and Edwards 1994).

The average $\delta^{13}\text{C}$ value of periphyton was -30.06‰. This was more depleted compared to other studies: -22.96‰ Puerto Rico (March and Pringle, 2003), -25.4‰ Hong Kong (Lau *et al.* 2009b) and -21‰ Guadeloupe (Coat *et al.* 2009). However, depleted $\delta^{13}\text{C}$ values of filamentous algae have been reported, such as -28.3‰ in Hong Kong (Mantel *et al.* 2004). The difference in periphyton $\delta^{13}\text{C}$ values reported in the literature compared to periphyton in our study could be a result of samples being taken from pools and riffles, biotopes with different environmental conditions compared to waterfalls. The $\delta^{13}\text{C}$ values of periphyton can be affected by several factors including the availability of dissolved inorganic carbon, taxonomic composition of the periphyton, water velocity, growth rate, and periphyton biomass (Hill and Middleton, 2006). The average $\delta^{15}\text{N}$ for periphyton (-1.29 ‰) was higher than in leaf litter (-0.91 ‰), which is not surprising as periphyton scrubs contain a complex mix of bacteria, algae, fungi, protozoa and can include animal tissue, thus increasing the $\delta^{15}\text{N}$ value (Wetzel, 2001; March and Pringle, 2003).

This study was conducted during a dry season when there are high rates of leaf litter fall (Cranbrook and Edwards, 1994; Pendry and Proctor, 1996). Leaves were therefore expected on all waterfalls during this study. However, during the rest of the year when there is less leaf litter fall and faster river flows, it would be predicted there would likely be less leaf litter. This study showed the importance of periphyton as a food source. Waterfalls are often situated under a canopy gap due to the wide plunge

Commented [BK2]: Reviewer comment: But does not leaf litter also include these extras??

The reviewer is right but periphyton is a biofilm that is made up of a mix of things, whereas leaf litter is mainly...leaf litter!

pools directly below the waterfall, which increase the channel width. Therefore, unlike other headwater tropical biotopes, waterfalls are less likely to be light limited. In addition to leaf litter and periphyton food sources, some macroinvertebrate guts contained fungal hyphae and conidia. It has long been acknowledged that microbial conditioning of leaf litter increases nutritional value and palatability for macroinvertebrates (Petersen and Cummins, 1974). In the tropics bacteria and fungi have been suggested to play a more important role in leaf litter decomposition due to the higher lignin content (Irons *et al.* 1994; Wallace *et al.* 1997; Boyero *et al.* 2015, 2016) and in the streams of Ulu Temburong it is not uncommon to see leaf litter covered in fungi.

Both Heptageniidae and Blephariceridae are defined as herbivorous grazers in the literature (Tonnoir 1930; Alverson *et al.* 2001; Ghee, 2004). However, in this study Blephariceridae had $\delta^{13}\text{C}$ values matching both periphyton and leaf litter. It is possible for the Blephariceridae to graze the periphyton and fungi that have colonized the leaf surface and as a consequence consume the leaf litter (Petersen and Cummins, 1974). The $\delta^{15}\text{N}$ values of Heptageniidae and Blephariceridae generally stayed under 2.5 ‰ implying these taxa are herbivorous (Table 2), with the GCA supporting these results.

In the literature Simuliidae and Hydropsychidae have been classed as filter feeders (Cummins, 1973; Cummins and Klug 1979; Fig. 3), however, these functional feeding group classifications are based on taxa living in temperate streams and there is a growing number of studies that suggest tropical taxa are more dependent on periphyton than leaf litter or detritus (March and Pringle, 2003; Brito *et al.* 2006; Lau *et al.* 2009a). For example, Brito *et al.* (2006) found Hydropsychidae in a Brazilian stream with $\delta^{13}\text{C}$ values matching aquatic macrophytes and microalgae. The low digestibility of tropical leaf litter as suggested by the C: N values in this study may

explain why periphyton is a more prevalent food source (Table 2; Benstead, 1996). Wootton (1977) argues that Simuliidae have been known to eat algae and will essentially ingest any particle from 0.5 μm to a maximum that depends on their mouth size (Burton, 1973). In addition, it is possible the water column contains many fragments of periphyton due to the frequent scouring flows. In Ulu Temburong National Park convective rainfall causes river and stream levels to fluctuate, this can occur daily in the wet season, with the power of the water scouring out habitats including periphyton (Sheldon, 2011). These consistent fast flows may have also influenced Simuliidae feeding behaviour causing them to scrape.

The SIA results showed that Simuliidae and Hydropsychidae had higher $\delta^{15}\text{N}$ values compared to herbivorous grazers (Heptageniidae and Blephariceridae; Table 2). The $\delta^{15}\text{N}$ values of consumers tend to be 2-5 ‰ higher than that of their diets (Hobson and Clark, 1992; Bearhop *et al.* 2002), therefore our results suggest that Simuliidae and Hydropsychidae on the waterfalls were omnivorous. Simuliidae have been known to consume Chironomidae (Serra-Tosio, 1967) and some studies have regarded Hydropsychidae as omnivorous and even predacious (Fuller and Mackay, 1980; Allan and Castillo, 1995). The GCA support the SIA with animal tissue present in the guts of both Simuliidae and Hydropsychidae. This study is the first known stable isotope analysis on Buccinidae snail, with the literature mainly focusing on its taxonomy and geographic distribution (Polgar *et al.* 2015). Our results show that the snail had little overlap with $\delta^{13}\text{C}$ values for leaf litter or periphyton (Table 2) and the $\delta^{15}\text{N}$ values confirm reports in the literature that Buccinidae is predacious (Coelho *et al.* 2013).

Some taxa showed high variability of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values (Table 2). This variability could be due to the coarse level of identification to family level. Identification to genus or species level may yield greater information, with species

potentially having different life histories and food preferences. However, this level of resolution is difficult in Borneo with macroinvertebrates still mostly undescribed (Baker *et al.* In press; Jacobsen *et al.* 2008; Yule and Yong, 2004). Variability in dietary composition could also be a result of substrate complexity along with food availability and quantity, which was not formally investigated. In our study the 'waterfalls' geomorphic complexity varied from smooth bedrock to substrate covered in numerous ridges and holes promoting the formation of a range of functional habitats defined by a mix of leaf litter, wood debris, moss and periphyton. Although these functional habitats were not quantitatively recorded in this study, visual observations found higher levels of wood debris and leaf litter on some waterfalls, particularly ones with complex substrate. Waterfall complexity may also influence abundance of different taxa on the waterfall and therefore number of prey for the predators, for example Buccinidae $\delta^{15}\text{N}$ value was lowest (4.35 ‰) at waterfall 9, which had smooth substrate. While, the highest level of $\delta^{15}\text{N}$ (6.63 ‰) was at waterfall 1 which had heterogeneous substrate. However, further studies are required to investigate if these changes in waterfall complexity influence macroinvertebrate feeding behaviour and trophic level.

While the food web is mostly resolved, it appears there are likely some unknown food sources. The first was a source with more negative $\delta^{13}\text{C}$ values (less than -33‰) that matched Blephariceridae and Heptageniidae at waterfall 10 (Fig. 3). Reviewing other tropical isotopic studies (including March and Pringle, 2003; Brito *et al.* 2006; Coat *et al.* 2009; Lau *et al.* 2009b) few food sources matched such low $\delta^{13}\text{C}$ values. However, Huang *et al.* (2007) did sample red algae, which had depleted ^{13}C with values around -37.4‰. Although this does not exactly match the $\delta^{13}\text{C}$ values of Blephariceridae and Heptageniidae it could suggest that the waterfalls do harbor some

types of algae with very depleted $\delta^{13}\text{C}$ values. The second 'unknown' food source had a $\delta^{13}\text{C}$ value of -27‰ and matched the $\delta^{13}\text{C}$ value for Heptageniidae at waterfall 7, Simuliidae at waterfall 11 and Hydropsychidae at waterfall 8 and 11. These values match values of dissolved organic matter and Fine Particulate organic matter (FPOM) from other studies, such as 28.6‰ on the Island of Guadeloupe (Coat *et al.* 2009) and in a Brazilian stream FPOM had a mean $\delta^{13}\text{C}$ value of $\sim 29\text{‰}$ (Brito *et al.* 2006).

Conclusions

Macroinvertebrate communities living on tropical waterfalls are significantly under researched, with this study being the first paper to examine their trophic structure. Results from both SIA and GCA show two, and potentially three, trophic levels occurring including the herbivore (Heptageniidae and Blephariceridae), omnivore (Simuliidae and Hydropsychidae) and predator (Buccinidae). In agreement with some tropical trophic structure studies on pools and riffles, the basal food resource on the waterfalls were found to be predominantly algae based. However, this study does suggest that despite the scouring velocities, waterfalls can retain functional habitats, with leaf litter being assimilated. The degree of functional habitat retention appears to be dependent on substrate heterogeneity and complexity. Quantitative analysis of substrate complexity is necessary to understand if the macroinvertebrate isotopic variation is related to waterfall complexity. In addition, further research is required to identify the unknown food sources.

Acknowledgements

This work was funded by a Natural Environment Research Council (NERC) funded PhD studentship (Code: DAR7132) and NERC Life Sciences Mass Spectrometry (Code: EK232-11/14). Thanks to the International Consortium of Universities for the Study of Biodiversity and the Environment (iCIBE) for fieldwork support and to NERC Life Sciences Mass Spectrometry Facility in East Kilbride for SIA support. Gut contents analysis was carried out with guidance from the Freshwater Biological Association River Labs.

REFERENCES

- Allan, J. D. and Castillo, M. M. (1995) *Stream ecology*. Springer.
- Alverson, A.J., Courtney, G.W. and Luttenton, M.R. (2001) Niche overlap of sympatric Blepharicera larvae (Diptera: Blephariceridae) from the southern Appalachian Mountains. *Journal of the North American Benthological Society* **20**, 564-581.
- Baker, K., Chadwick, M.A., Wahab, R.A. and Kahar, R. (2016) Benthic community structure and ecosystem functions in above-and below-waterfall pools in Borneo. *Hydrobiologia*, 1-16.
- Baker, K., Chadwick, M. A. Wahab, R. A. Kahar R. and Sulaiman, Z. in press. Fluvial biotopes influence macroinvertebrate biodiversity in Southeast Asian tropical streams. *Ecosphere*.
- Bearhop, S., Adams, C. E., Waldron, S., Fuller, R. A. and Macleod, H. (2004) Determining trophic niche width: a novel approach using stable isotope analysis. *Journal of Animal Ecology* **73**, 1007-1012.
- Benstead, J. P. (1996) Macroinvertebrates and the Processing of Leaf Litter in a Tropical Stream. *Biotropica* **28**, 367-375.
- Brito, E., Moulton, T., De Souza, M. and Bunn, S. (2006) Stable isotope analysis indicates microalgae as the predominant food source of fauna in a coastal forest stream, south-east Brazil. *Austral ecology* **31**, 623-633.
- Boyero, L., Pearson, R.G., Hui, C., Gessner, M.O., Pérez, J., Alexandrou, M.A., Graça, M.A., Cardinale, B.J., Albariño, R.J., Arunachalam, M. and Barmuta, L.A. (2016) Biotic and abiotic variables influencing plant litter breakdown in streams: a global study. In *Proc. R. Soc. B* (Vol. 283, No. 1829, p. 20152664). The Royal Society.

492 Boyero, L., Pearson, R.G., Gessner, M.O., Dudgeon, D., Ramírez, A., Yule, C.M., Callisto, M.,
 493 Pringle, C.M., Encalada, A.C., Arunachalam, M. and Mathooko, J., 2015. Leaf-litter
 494 breakdown in tropical streams: is variability the norm?. *Freshwater Science*, 34(2), pp.759-
 495 769.

496 Burton, G. J. (1973) Feeding of *Simulium Hargreavesi* Gibbins Larvae on *Oedogonium* Algal
 497 Filaments in Ghana. *Journal of Medical Entomology* **10**, 101-106.

498 Clayton, P. D. and Pearson, R. G. (2016) Harsh habitats? Waterfalls and their faunal dynamics in
 499 tropical Australia. *Hydrobiologia* **775**, 123-137.

500 Coat, S., Monti, D., Bouchon, C. and Lepoint, G. (2009) Trophic relationships in a tropical stream food
 501 web assessed by stable isotope analysis. *Freshwater Biology* **54**, 1028-1041.

502 Coelho, A. R., Dinis, M. T. and Reis, J. (2013) Effect of diet and stocking densities on life history traits
 503 of *Clea helena* (Philippi 1847) reared in captivity. *Journal of Aquaculture Research &*
 504 *Development* **187**.

505 Coughlan, J. F., Pearson, R. G. and Boyero, L. (2010) Crayfish process leaf litter in tropical streams
 506 even when shredding insects are common. *Marine and Freshwater Research* **61**, 541-548.

507 Covich, A. P., Crowl, T. A., Hein, C. L., Townsend, M. J. and McDowell, W. H. (2009) Predator-prey
 508 interactions in river networks: comparing shrimp spatial refugia in two drainage basins.
 509 *Freshwater Biology* **54**, 450-465.

510 Cranbrook, Earl of., and Edwards, D.S (1994) *Belalong a tropical rainforest*. (Sun Tree Publishing Ltd:
 511 Singapore.)

512 Crowl, T. A., McDowell, W. H., Covich, A. P. and Johnson, S. L. (2001) Freshwater shrimp effects on
 513 detrital processing and nutrients in a tropical headwater stream. *Ecology* **82**, 775-783.

514 Cummins, K. W. (1973) Trophic relations of aquatic insects. *Annual review of entomology* **18**, 183-
 515 206.

516 Cummins, K. W. and Klug, M. J. (1979) Feeding ecology of stream invertebrates. *Annual review of*
 517 *ecology and systematics* **10**, 147-172.

518 Dykes, A. P. (1994) Landform Processes. In: Cranbrook, Earl of. & Edwards D. S. (Eds) *Belalong, a*
 519 *tropical rainforest*. Sun Tree Publishing Ltd, Singapore.

520 Dykes, A. P. (1996) Analysis of factors contributing to the stability of steep hillslopes in the tropical
 521 rainforest of Temburong, Brunei Darussalam. In: D. S. Edwards, W. E. Booth & S. C. Choy

522 Eds. *Tropical Rainforest Research — Current Issues*. Vol. 74: Springer Netherlands, 1996,
523 387-409. Edwards, D. S. (1994) *Belalong, a tropical rainforest*. Singapore: Sun Tree
524 Publishing.

525 Flecker, A. S. and Taylor, B. W. (2004) Tropical Fishes as Biological Bulldozers: Density Effects on
526 Resource Heterogeneity and Species Diversity. *Ecology* **85**, 2267-2278.

527 Fry, B. (1991) Stable Isotope Diagrams of Freshwater Food Webs. *Ecology* **72**, 2293-2297.

528 Fry, B. (2007) Stable isotope ecology. Springer Science & Business Media, United States.

529 Fuller, R. L. and Mackay, R. J. (1981) Effects of food quality on the growth of three Hydropsyche
530 species (Trichoptera: Hydropsychidae). *Canadian Journal of Zoology* **59**, 1133-1140.

531 Furtado, J. I. (1969) Ecology of Malaysian odonates: Biotope and association of species. *Verh.*
532 *Internat. Verein. Limnol* **17**, 863-887.

533 Ghee, K. S. (2004) Insecta: Ephemeroptera. In: C. M. Yule & H.-S. Yong Eds. *Freshwater*
534 *invertebrates of the Malaysian region*. Malaysia: Academy of Sciences Malaysia Kuala
535 Lumpur, 2004.

536 Giller, P.S., and Malmqvist, B.. (1998) The biology of streams and rivers (Oxford University Press:
537 Oxford.)

538 Grey J, Jones, R. I. and Sleep, D. (2001) Seasonal changes in the importance of the source of organic
539 matter to the diet of zooplankton in Loch Ness, as indicated by stable isotope analysis. *Limnol*
540 *Oceanogr* **46**, 505-513

541 Harper, D. M., Smith, C. D., Barham, P. J. and Howell, R. (1995) The ecological basis for the
542 management of the natural river environment. In: D. M. Harper & A. J. D. Ferguson Eds. *The*
543 *Ecological Basis for River Management*. Chichester: John Wiley, 1995, 219-238.

544 Hart, D. D. and Finelli, C. M. (1999) Physical-biological coupling in streams: the pervasive effects of
545 flow on benthic organisms. *Annual Review of Ecology and Systematics* **30**, 363-395.

546 Hein, C. L. and Crowl, T. A. (2010) Running the predator gauntlet: do freshwater shrimp (*Atya*
547 *lanipes*) migrate above waterfalls to avoid fish predation? *Journal of the North American*
548 *Benthological Society* **29**, 431-443.

549 Hershey, A. E. and Peterson, B. J. (1996) Stream food webs. In: H. F.R. & L. G.A. Eds. *Methods in*
550 *Stream Ecology* London: Academic, 1996, 511-530.

551 Hill, W. R. and Middleton, R. G. (2006) Changes in carbon stable isotope ratios during periphyton
552 development. *Limnology and Oceanography* **51**, 2360-2369.

553 Hobson, K. A. and Clark, R. G. (1992) Assessing Avian Diets Using Stable Isotopes I: Turnover of ^{13}C
554 in Tissues. *The Condor* **94**, 181-188.

555 Huang, I., Lin, Y., Chen, C. and Hsieh, H. (2007) Food web structure of a subtropical headwater
556 stream. *Marine and Freshwater Research* **58**, 596-607.

557 Hury, A.D., Riley, R.H., Young, R.G., Arbuckle, C.J. and Peacock, K. (2002). Natural-abundance
558 stable C and N isotopes indicate weak upstream-downstream linkage of food webs in a
559 grassland river. *Archiv für Hydrobiologie* **153**, 177-196.

560 Irons, J. G., Oswood, M. W., Stout, R. J. and Pringle, C. M. (1994) Latitudinal patterns in leaf litter
561 breakdown: is temperature really important? *Freshwater Biology* **32**, 401-411.

562 Jacobsen, D., C. Cressa, J. M. Mathooko, and D. Dudgeon. 2008. Macroinvertebrates: Composition,
563 Life Histories and Production. Pages 65-105 in D. Dudgeon, editor. *Tropical Stream Ecology*.
564 Academic Press, London.

565 Kano, Y., Nishida, S. and Nakajima, J. (2012) Waterfalls drive parallel evolution in a freshwater goby.
566 *Ecology and Evolution* **2**, 1805-1817.

567 Lau, D. C. P., Leung, K. M. Y. and Dudgeon, D. (2009) What does stable isotope analysis reveal about
568 trophic relationships and the relative importance of allochthonous and autochthonous
569 resources in tropical streams? A synthetic study from Hong Kong. *Freshwater Biology* **54**,
570 127-141.

571 Layer, K., Hildrew, A. G. and Woodward, G. (2012) Grazing and detritivory in 20 stream food webs
572 across a broad pH gradient. *Oecologia* **171**, 459-471.

573 Li, A. O. Y. and Dudgeon, D. (2008) Food resources of shredders and other benthic macroinvertebrates
574 in relation to shading conditions in tropical Hong Kong streams. *Freshwater Biology* **53**,
575 2011-2025.

576 Mantel, S. K., Salas, M. and Dudgeon, D. (2004) Foodweb structure in a tropical Asian forest stream.
577 *Journal of the North American Benthological Society* **23**, 728-755.

578 March, J. G., Pringle, C. M., Townsend, M. J. and Wilson, A. I. (2002) Effects of freshwater shrimp
579 assemblages on benthic communities along an altitudinal gradient of a tropical island stream.
580 *Freshwater Biology* **47**, 377-390.

581 March, J. G. and Pringle, C. M. (2003) Food Web Structure and Basal Resource Utilization along a
582 Tropical Island Stream Continuum, Puerto Rico. *Biotropica* **35**, 84-93.

583 Newson, M. and Newson, C. (2000) Geomorphology, ecology and river channel habitat: mesoscale
584 approaches to basin-scale challenges. *Progress in Physical Geography*, 24(2), 195-217.

585 Odland, A., Birks, H. H., Botnen, A., Tønsberg, T. and Vevle, O. (1991) Vegetation change in the
586 spray zone of a waterfall following river regulation in Aurland, western Norway. *Regulated
587 Rivers: Research & Management* **6**, 147-162.

588 Pendry, C. A. and Proctor, J. (1996) The Causes of Altitudinal Zonation of Rain Forests on Bukit
589 Belalong, Brunei. *Journal of Ecology* **84**, 407-418.

590 Petersen, R. C. and Cummins, K. W. (1974) Leaf processing in a woodland stream. *Freshwater biology*
591 **4**, 343-368.

592 Peterson, B. J. and Fry, B. (1987) Stable isotopes in ecosystem studies. *Annual review of ecology and
593 systematics* **18**, 293-320.

594 Polgar, G., Khang, T. F., Chua, T. and Marshall, D. J. (2015) Gross mismatch between thermal
595 tolerances and environmental temperatures in a tropical freshwater snail: Climate warming
596 and evolutionary implications. *Journal of Thermal Biology* **47**, 99-108.

597 Pringle, C. M. and Hamazaki, T. (1998) The role of omnivory in a neotropical stream: separating
598 diurnal and nocturnal effects. *Ecology* **79**, 269-280.

599 Rackemann, S. L., Robson, B. J. and Matthews, T. G. (2013) Conservation value of waterfalls as
600 habitat for lotic insects of western Victoria, Australia. *Aquatic Conservation: Marine and
601 Freshwater Ecosystems* **23**, 171-178.

602 Ramirez, A. and Hernandez-Cruz, L. R. (2004) Aquatic Insect Assemblages in Shrimp-dominated
603 Tropical Streams, Puerto Rico. *Biotropica* **36**, 259-266.

604 Rosi-Marshall, E. J. and Wallace, J. B. (2002) Invertebrate food webs along a stream resource gradient.
605 *Freshwater Biology* **47**, 129-141.

606 Serra-Tosio, B., 1967. La prise de nourriture chez la larve de *Prosimulium inflatum* Davies, 1957
607 (Diptera, Simuliidae). *Travaux du Laboratoire d'hydrobiologie et de pisciculture de
608 l'Université de Grenoble*, pp.57-58. From: Wotton, R. S. (1977) The Size of Particles
609 Ingested by Moorland Stream Blackfly Larvae (Simuliidae). *Oikos* **29**, 332-335.

Sheldon, A. L. (2011) Comparative habitat use by grazing fishes in a Bornean stream. *Environmental Biology of Fishes* **92**, 381-390.

Thomas, R., Vaughan, I. and Lello, J. (2013) *Data analysis with R statistical software: a guidebook for scientists*. Eco-explore, Cardiff.

Tonnoir, A. (1930) Notes on Indian Blepharocid larvae and pupae with remarks on the morphology of Blepharocid larvae and pupae in general. *Records of the Indian Museum* **32**, 161-214.

Torrente-Vilara, G., Zuanon, J., Leprieur, F., Oberdorff, T. and Tedesco, P. A. (2011) Effects of natural rapids and waterfalls on fish assemblage structure in the Madeira River (Amazon Basin). *Ecology of Freshwater Fish* **20**, 588-597.

Townsend, C. R. and Hildrew, A. G. (1994) Species traits in relation to a habitat templet for river systems. *Freshwater Biology* **31**, 265-275.

Wallace, J. B., Eggert, S. L., Meyer, J. L. and Webster, J. R. (1997) Multiple trophic levels of a forest stream linked to terrestrial litter inputs. *Science* **277**, 102-104.

Wetzel, R. G. (2001) *Limnology: lake and river ecosystems, 3rd edn*. San Diego: Academic Press.

Wotton, R. S. (1977) The Size of Particles Ingested by Moorland Stream Blackfly Larvae (Simuliidae). *Oikos* **29**, 332-335.

Yule, C. M., Boyero, L. and Marchant, R. (2010) Effects of sediment pollution on food webs in a tropical river (Borneo, Indonesia). *Marine and Freshwater Research* **61**, 204-213.

Yule, C. M., and Yong, H. S. (2004). 'Freshwater Invertebrates of the Malaysian Region'. (Academy of Sciences Malaysia: Kuala Lumpur.)

Tables

Table 1. Physical measurements of waterfalls including average slope angle and length. Presence (1) vs. absence (0) of functional habitats (leaf litter, wood debris and moss).

Waterfall	Slope angle (°)	Length (m)	Leaf litter	Wood Debris	Moss
1	53	5.7	1	1	0
2	26	4.93	1	0	0

3	20	6.16	1	0	0
4	50	15	1	1	1
5	47	7.2	1	0	1
6	39	7	1	0	1
7	47	~6	1	1	1
8	35	20.4	1	0	0
9	27	~13	1	0	0
10	16	9.13	1	0	0
11	40	4.47	1	0	1
12	48	16.2	1	1	0

Table 2. Average carbon and nitrogen isotopic values of basal food sources and consumers (mean \pm SD).

	<i>n</i>	C:N	$\delta^{13}\text{C}$ (‰): Mean \pm SD	$\delta^{15}\text{N}$ (‰): Mean \pm SD
Basal food sources				
Leaf litter	36	49.27	-32.97 \pm 0.26	-0.91 \pm 0.16
Periphyton	57	5.37	-30.06 \pm 0.34	1.29 \pm 0.10
Consumers				
Heptageniidae	36	4.56	-29.47 \pm 0.33	1.06 \pm 0.13
Blephariceridae	30	4.50	-31.45 \pm 0.42	1.50 \pm 0.12
Simuliidae	36	4.30	-28.14 \pm 0.09	2.16 \pm 0.06
Hydropsychidae	36	4.53	-28.58 \pm 0.14	3.07 \pm 0.08
Buccinidae	33	3.92	-26.18 \pm 0.16	5.48 \pm 0.12

Figure captions

Fig. 1. Map showing the location of Brunei Darussalam in northern Borneo (A). Brunei Darussalam is split into two parts with this study being conducted in the Temburong district, the eastern section of Brunei (B). The twelve waterfalls are situated on tributaries of the two main rivers Sungai Belalong and Sungai Temburong (C) near to Kuala Belalong Field Study Centre (KBFSC)

Fig. 2. Average percent contributions of stomach contents including amorphous detritus, animal tissue, CPOM, algae and fungal hyphae and conidia in the five macroinvertebrate families - Hydropsychidae, Heptageniidae, Blephariceridae and Simuliidae.

Fig. 3. Mean $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of basal resources leaf litter (LL) and periphyton (P) for the abundant macroinvertebrates found on the waterfalls in Ulu Temburong National Park. Macroinvertebrates include Buccinidae, Blephariceridae, Heptageniidae, Hydropsychidae and Simuliidae. Error bars indicate ± 1 SD.

Fig. 4. Food web of the abundant macroinvertebrates present on waterfalls in Ulu Temburong National Park. Based on stomach contents analysis and stable isotope analysis. Line thickness represents contributions of the sources.

Accepted Draft (2017)